



A note on definition of matrix convex functions[☆]

Oleg E. Tikhonov

Research Institute of Mathematics and Mechanics, Kazan State University, Nuzhina 17, Kazan 420008, Russia

Received 6 May 2005; accepted 20 December 2005

Available online 8 February 2006

Submitted by C. Davis

Abstract

We prove that a real-valued function f defined on an interval S in \mathbf{R} is matrix convex if and only if for any natural k , for all families of positive operators $\{A_i\}_{i=1}^k$ in a finite-dimensional Hilbert space, such that $\sum_{i=1}^k A_i = 1$, and arbitrary numbers $x_i \in S$, the inequality

$$f\left(\sum_{i=1}^k x_i A_i\right) \leq \sum_{i=1}^k f(x_i) A_i$$

holds true.

© 2006 Elsevier Inc. All rights reserved.

Keywords: Matrix convex function; The Neumark theorem

The aim of this note is to add one more characterization to the list of known characterizations of matrix and operator convex functions (see, e. g., [1,3,4]).

In what follows, S stands for an interval in \mathbf{R} of an arbitrary type. For a function $f: S \rightarrow \mathbf{R}$ and a self-adjoint operator X in a finite-dimensional Hilbert space H with spectrum in S , the value $f(X)$ is defined by the spectral theorem. The identity operator in H is denoted by 1_H .

Theorem. For a function $f: S \rightarrow \mathbf{R}$, the following conditions are equivalent:

(i) f is matrix convex, i.e.,

$$f(\alpha X + (1 - \alpha)Y) \leq \alpha f(X) + (1 - \alpha)f(Y)$$

[☆] Supported by the Russian Foundation for Basic Research (grant no. 05-01-00799).

E-mail address: Oleg.Tikhonov@ksu.ru